Saranthan et al.

S/N: 09/681,068

In the Claims

1. (Original) A method of acquiring free-breathing MR images comprising the steps of:

monitoring heart rate of a subject just prior to image acquisition to acquire a time period of an R-R interval;

recording the time period from the heart rate monitoring to prospectively estimate future R-R intervals; and

acquiring n sets of MR data, a first MR data acquisition commencing at any point in an R-R interval and extending for the time period recorded.

- 2. (Original) The method of claim 1 further comprising the steps of segmenting each MR data acquisition into n segments and repetitively acquiring each segment in n successive heartbeats.
- 3. (Original) The method of claim 2 further comprising the step of combining the n MR data sets to form a set of MR images with high temporal resolution covering the R-R interval.
- 4. (Original) The method of claim 1 further comprising the step of discontinuing heart rate monitoring before acquiring MR image data.
- 5. (Original) The method of claim 1 wherein a second set of MR data is acquired immediately after the acquisition of the first set of MR data.
 - 6. (Original) The method of claim 2 wherein n=1 for fluoroscopy imaging.
- 7. (Original) The method of claim 1 wherein the step of acquiring MR data is performed using one of a fast gradient-recalled echo pulse sequence and a steady state free precession pulse sequence.
- 8. (Original) The method of claim 1 further comprising the steps of:
 subjecting a patient to successively increased, graded levels of cardiac stress
 during the monitoring step until the heart rate is stabilized at a required stress level; and

BEST AVAILABLE COPY

Saranthan et al.

S/N: 09/681,068

acquiring MR data according to the acquisition step of several long and short axis views of at least a portion of a heart muscle to monitor cardiac function during any portion of a stress test.

- 9. (Original) The method of claim 8 wherein the cardiac stress is induced either by physical stress or administration of a pharmaceutical.
- 10. (Withdrawn) A computer program for use with an MRI scanner having a computer, the computer program having a set of instruction that, when executed, cause the computer to:

receive a time-period signal indicative of an R-R interval representing a cardiac cycle of a patient;

acquire a first set of partial MR image data during a first acquisition period equal to the R-R interval;

All of the claims are not present.
Colled

Jamara